

ORES11.001APC

PATENT

**POLYMER STABILISED, CRYSTALLISED, CATANIONIC MEMBRANES,
PREPARATION METHOD THEREOF AND APPLICATIONS OF SAME**

The present invention relates to polymer-stabilized crystalline catanionic membranes, to the
5 process for preparing them and to the uses thereof, especially as medicaments for vectorizing active species or for retaining, via adsorption, volatile molecules.

Mixtures of anionic and cationic surfactants
10 in aqueous medium give rise to what may conveniently be referred to as "catanionic" solutions.

After ionic pairing, the counterions form a salt in excess and induce high conductivity of the samples that masks the electrostatic interactions. One
15 particular type of salt-free catanionic formulation is obtained by using only H^+ and OH^- counterions, such that no excess of salt is formed by the mixing of the two surfactants (Dubois M. et al., C. R. Acad. Sci. Paris II C, 1998, 1(9) 567-565). The resulting
20 catanionic systems are commonly known as "true catanionic systems".

When these catanionic solutions are heated to a temperature above the melting point of the chains, the anionic and cationic surfactants self-assemble in
25 the form of stable micelles of varied forms (spheres, cylinders or folded bilayers). Depending on the relative proportions between cationic and anionic constituents, various forms of structures may then be obtained during the cooling of these solutions.

When the catanionic solution contains an excess of cationic surfactants, the formation of crystalline nanodisks is observed, which are formed from a sandwich structure with a rigid outer membrane, the adjustable diameter of which may range from a few
35 microns to about thirty nanometers and in which the positive charges are mainly located in the slice. The structure and the process for preparing these catanionic nanodisks are especially described in the article by Zemb T. et al., Science, 1999, **283**, 816-819.

Conversely, when the catanionic solution contains an excess of anionic surfactants, the formation of hollow polyhedra is observed, the shape of which will vary as a function of the amount of anionic surfactants in excess. Under certain conditions, the formation of hollow icosahedra is observed in particular, the shape of which is reminiscent of that observed for viral capsid proteins. Their structure and the process for preparing them are especially described in the article by Dubois M. et al., Nature, 2001, **411**, 672-675. Said article especially describes micrometer-sized icosahedra weighing about 10^{10} daltons, the structure of which is partly stabilized by the presence of pores at the apices of the polyhedra. According to said article, the formation of an icosahedron requires the conjunction of the following three conditions:

- 1) the formation of stable unilamellar vesicles during the high-temperature equimolar mixing of the anionic and cationic surfactant solutions,
- 2) the excess of anionic surfactant must be insoluble in water and in the crystalline bilayer obtained during the equimolar mixing of the anionic and cationic surfactants,
- 3) the amount of surfactant in excess must be such that it allows the formation of 10 to 15 pores per vesicle.

The absence of condition 3) leads to the formation of open crystalline large bilayers or nanodisks including pores.

Still according to the teaching of said article and by virtue of the presence of these pores, various uses of such polyhedra may be envisioned. They may be used, for example, as medicaments for the controlled release of active principles or of DNA in gene therapy or the isolation of RNA strands in order to protect them against the action of destructive enzymes. However, these polyhedra have the major drawback of being particularly sensitive to the slightest presence of salts, and have a tendency to

aggregate together, which prohibits, for example, their use in physiological media, for instance in blood, and consequently their intravenous injection.

Moreover, various structures for delivering
5 active principles, including sustained-release forms, have already been proposed and are based on the encapsulation of active ingredients inside vesicles. Thus, with this aim, a large number of prior art documents have described the use of spherical vesicles
10 consisting of one or more lipid bilayers commonly denoted as liposomes. However, the use of liposomes is not always entirely satisfactory in terms of stability especially, and also since their preparation involves processes that require the use of organic solvents, the
15 use of which is not necessarily compatible with physiological media and a particular material.

Thus, in order to overcome all these drawbacks and to provide a novel system for delivering active molecules that is stable, resistant to high
20 ionic strengths and simple to use, the Inventors have developed that which forms the subject of the present invention.


A first subject of the invention is thus a catanionic membrane in the form of an organized solid
25 bilayer comprising a lateral alternation of anionic surfactants with H^+ counterions and of cationic surfactants with cocrystallized OH^- counterions in which the mole fraction (MF): molar amount of anionic surfactants (Q_{AS})/(molar amount of anionic surfactants
30 (Q_{AS}) + molar amount of cationic surfactants (Q_{CS})) is greater than 0.5 (i.e. $Q_{AS}/(Q_{AS} + Q_{CS}) > 0.5$), said membrane forming a surface that is at least locally flat, characterized in that said bilayer is stabilized with at least one polymer that is neutral and hydro-
35 phobic or of overall electrical charge opposite the effective charge of said catanionic membrane, said polymer being adsorbed onto said surface.

The presence of the polymers absorbed onto their surface allows the catanionic membranes in

accordance with the invention to be stabilized during Brownian motion, and in particular allows a distance of at least one nanometer to be conserved between two membranes, avoiding their precipitation and thus
5 allowing their dilution with isotonic solutions such as seawater or blood.

The cationic and anionic surfactants that may be used in accordance with the invention are preferably chosen from compounds with a melting point greater than
10 the working temperature so as to be in crystalline form. Since the working temperature may range between 20 and 30°C, surfactants with a melting point of greater than 30°C will more particularly be selected.

The cationic surfactants that may be used in accordance with the invention for the formation of the bilayers are preferably chosen from the monocationary and bicationary quaternary ammoniums of formulae (I) and (I'), respectively, below:



in which:

- R₁, R₂ and R₃, which may be identical or different, represent a C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl
25 or (C₁-C₄)alkyl ether radical,

- R'₁ and R'₂, which may be identical or different, represent a C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl or (C₁-C₄)alkyl ether radical,

- R'₃ and R'₄, which may be identical or different, represent a saturated or unsaturated C₈-C₂₄
30 hydrocarbon-based chain, a benzyl or (C₄-C₂₀)alkylbenzyl radical or a (C₄-C₂₀)alkyl ester group,

- R₄ represents a saturated or unsaturated C₈-C₂₄ hydrocarbon-based chain, a benzyl or
35 (C₄-C₂₀)alkylbenzyl radical or a (C₄-C₂₀)alkyl ester group;
and mixtures thereof.

Among the C_1 - C_4 alkyl radicals of the cationic surfactants of formulae (I) and (I') above, the methyl radical is particularly preferred.

5 Among the C_8 - C_{24} hydrocarbon-based chains of the radicals R'_3 , R'_4 and R'_4 , mention may be made especially of alkyl chains, for instance stearyl, cetyl, dodecyl and tetradecyl chains.

Among the (C_4 - C_{20})alkyl ester groups mentioned for the radicals R'_3 , R'_4 and R_4 , mention may be made in particular of (C_{16})alkyl esters and (C_{12})alkyl esters.

10 Among the compounds of formula (I) above with OH^- counterions, mention may be made in particular of cetyltrimethylammonium hydroxide, dodecyltrimethylammonium hydroxide, stearyltrimethylammonium hydroxide, 15 tetradecyltrimethylammonium hydroxide, N-(2-carboxyethyl)-N,N-dimethyl-1-hexadecanaminium hydroxide, N-(2-hydroxyethyl)-N,N-dimethyl-1-hexadecanaminium hydroxide, cetyltriethylammonium hydroxide, dodecyltriethylammonium hydroxide, stearyltriethylammonium hydroxide, tetradecyltriethylammonium hydroxide, cetyltripropylammonium hydroxide, dodecyltripropylammonium hydroxide, stearyltripropylammonium hydroxide and tetradecyltripropylammonium hydroxide.

20 Among the compounds of formula (I') above, mention may be made in particular of didodecyldimethylammonium hydroxide, didodecyldiethylammonium hydroxide, didodecyldipropylammonium hydroxide, didodecyldibutylammonium hydroxide and dicetyldimethyltrimethylammonium hydroxide.

30 The anionic surfactants that may be used in accordance with the invention for the formation of the bilayers are preferably chosen from carboxylic acids with a C_8 - C_{24} carbon-based hydrophobic chain with H^+ counterions and phosphates and sulfonates with H^+ counterions comprising one or two C_{12} - C_{20} alkyl chains.

35 Among these anionic surfactants, mention may be made in particular of fatty acids such as myristic acid, lauric acid and palmitic acid, phosphates, sulfonates, benzyl sulfates and monocatenary glycerol

monoesters, preferably having a fine X-ray diffraction peak at large angles after combination with the cationic surfactant, located at $q = 1.52 \pm 0.15 \text{ \AA}^{-1}$ without a shoulder, as described, for example, by
5 Rank J.L. et al., J. Mol. Biol., 1974, **85**(2), 249-277.

It is possible, in accordance with the invention, to combine any type of anionic surfactant with H^+ counterions with any type of cationic surfactant with OH^- counterions and in particular with
10 any cationic surfactant of formula (I) and/or of formula (I') since it is possible to envision mixing of monocatenary and bicatenary cationic surfactants.

As has been indicated previously, one of the essential characteristics of the invention is that the
15 MF of the surfactants used for the formation of the bilayers ($Q_{AS}/(Q_{AS} + Q_{CS})$) should be greater than 0.5. According to one preferred embodiment of the invention, this MF is between 0.52 and 0.66 and even more preferentially between 0.55 and 0.58.

20 According to one particular and preferred embodiment of the invention, the bilayers consist:

a) either of a cationic surfactant of formula (I) as defined above and in which the radicals R_1 , R_2 and R_3 are identical and represent a methyl radical and
25 R_4 represents a hydrocarbon-based chain containing X carbon atoms, X being between 8 and 24 inclusive, combined with a carboxylic acid as defined above in which the C_8-C_{24} carbon-based hydrophobic chain contains $X \pm 4$ carbon atoms;

30 b) or of a cationic surfactant of formula (I') as defined above in which the radicals R'_1 and R'_2 are identical and represent a methyl radical and R'_3 and R'_4 are identical and represent a hydrocarbon-based chain containing X carbon atoms, X being between 8 and
35 24 inclusive, combined with a carboxylic acid as defined above in which the C_8-C_{24} carbon-based hydrophobic chain contains $X \pm 4$ carbon atoms;

c) or a phosphate or a sulfonate comprising two identical alkyl chains containing X carbon atoms, X

being between 8 and 24 inclusive, combined with a cationic surfactant of formula (I) as defined above and in which the radicals R_1 , R_2 and R_3 are identical and represent a methyl radical and R_4 represents a C_8 - C_{24} alkyl chain;

d) or a phosphate or a sulfonate comprising only one alkyl chain containing X carbon atoms, X being between 8 and 24 inclusive, combined with a cationic surfactant of formula (I') as defined above and in which the radicals R'_1 and R'_2 are identical and represent a methyl radical and R'_3 and R'_4 are identical and represent a C_8 - C_{24} alkyl chain.

According to an even more particularly preferred form of the invention, the bilayers are formed from a combination of cetyltrimethylammonium with OH^- counterions and of myristic acid with H^+ counterions.

In addition to the anionic surfactants with H^+ counterions, the bilayers in accordance with the invention may also contain a minor molar amount of anionic surfactants with metal counterions, and in particular with a sodium, magnesium, lithium, chromium, vanadium or nickel counterion, for example, said surfactants being chosen from the anionic surfactants mentioned above, with the exception, of course, of the nature of the counterion. According to the invention, the term "minor" molar amount means an amount of anionic surfactant with a metallic counterion that is necessary to bring the final MF within the range of between 0.52 and 0.66 as defined above.

According to one particular embodiment of the invention, the pairs of ions forming the polar heads of the cationic and anionic surfactants perpendicularly occupy an area preferably equal to that of the crystalline chains, on a hexagonal network, which may be detected by a first fine peak located at about $q = 1.52 \pm 0.15 \text{ \AA}^{-1}$.

According to one advantageous embodiment of the invention, the polymers that may be used to

stabilize the bilayers in accordance with the invention are nonlipid polymers, chosen from polymers that are neutral or that have an overall electrical charge opposite the effective charge of the catanionic
5 membranes, i.e. of "weakly" negative overall electrical charge.

For the purposes of the present invention, the term "effective charge" means the overall electrical charge taking into account the apparent pKa
10 of the acid in the crystal, which is manifested, for example, in electrophoretic mobility by a movement of the bilayers towards the anode. This electrical charge is different than the assayable structural charge and deduced from the localization of the composition in a
15 phase diagram.

Also for the purposes of the present invention, the term "polymers of "weakly" negative overall electrical charge means polymers comprising less than one elemental electrical charge per 2 nano-
20 meters of drawn length.

When they are neutral polymers, they are preferably chosen from polysaccharides, for instance dextrans and cellulose derivatives such as hydroxymethylcelluloses, hydroxyethylcelluloses and hydroxypropylcelluloses, synthetic polymers such as
25 polyethylene glycols (PEG), polyoxyethylenes, polyvinylpyrrolidone (PVP) and polyvinyl alcohols such as the products sold under the trade names PVA, Ethenol®, Poval®, Acroflex®, Airvol®, Alcotex® or Aquafilm®,
30 oxyethylenated diblock polymers such as the polymers sold under the trade name Varonic® by the company Degussa-Goldschmidt, block copolymers based on ethylene oxide and propylene oxide, such as the polymers sold under the trade names Pluronic® and Lutrol® by the
35 company BASF, and water-soluble equivalents thereof, water-soluble triblock copolymers, i.e. copolymers composed of hydrophilic-hydrophobic-hydrophilic blocks, such as the products sold under the trade names Methyl oxirane, EOPO copolymer, Antarox®, Arcol®, Daltocel® or

Dowfax® and analogs thereof comprising polystyrene as hydrophobic group.

When they are polymers of weakly negative overall electrical charge, they are preferably chosen
5 from polyacrylates, polymethacrylates, polyethyl methacrylates, polybutyl methacrylates and polystyrene-sulfonates, said polymers being substituted to more than 75% randomly with neutral water-soluble groups, for instance polyoxyethylene groups or the like.

10 Among the polymers that may be used in accordance with the invention, it is most particularly preferred to use weakly adsorbed polymers such as polyoxyethylene, dextran, PVP or the polymers sold under the trade names Varonic®, Pluronic® and Lutrol®,
15 Methyl Oxirane, Pluronic®, AntaroX®, Arcol®, Daltocel® and Dowfax®.

Among these polymers, it is even more particularly preferred to use a polyethylene glycol with a molecular mass of between 5000 and 50 000 Da and
20 even more preferentially between 10 000 and 20 000 Da.

These polymers preferably represent from 10% to 400% and even more particularly from 100% to 200% by weight relative to the total weight of the bilayer.

According to a particular embodiment of the
25 invention, the catanionic membranes may especially be in the form of faceted hollow microcrystals when the $\frac{MF_{QAS}}{Q_{AS} + Q_{CS}}$ is between 0.55 and 0.58.

In this case, these microcrystals may take the form of hollow polyhedra (molecular boxes)
30 comprising from 12 to 30 approximately triangular faces, and most particularly in the form of hollow icosahedra with an inner volume of between 0.1 and 10 μ^3 .

Within the organized solid bilayer of each of
35 the faces of these microcrystals, the lateral alternation of the cocrystallized anionic and cationic surfactants is hexagonal, the flat part of said faces consisting solely of species containing H^+ or OH^- counterions in stoichiometric amounts, whereas the

apices of said faces are in the form of an internal semitorus predominantly formed from the anionic species in excess and in an amount sufficient to obtain an MF $Q_{As}/(Q_{As} + Q_{Cs})$ of between 0.55 and 0.58.

5 According to this configuration, i.e. when the apices of each face are in the form of an internal semitorus, then the apex of each of the faces of a microcrystal forms a pore, together with the apices of the adjacent faces of the same microcrystal. In this
10 case, each microcrystal may comprise from about 10 to 15 pores.

By virtue of the presence of the neutral polymers or polymers with a weakly negative electrical charge adsorbed onto their surface, which block
15 flocculation and coalescence, these molecular boxes can withstand the ionic strength, i.e. the presence of salts, up to the point of isotonicity. The presentation of the catanionic membranes in molecular-box form is particularly preferred according to the invention.

20 The catanionic membranes in accordance with the invention may also be in the form of fragments of hollow polyhedra, i.e. in the form of a stack of three-dimensional catanionic crystals in the form of a "pile of plates", resulting from the opening of the molecular
25 boxes and the dense stacking of the fragments of the facets.

After their formation and before the adsorption of the specific stabilizing polymer, the pH of the membrane solution may be adjusted to any pH
30 value of between 2 and 6. This allows reactions to be performed in acidic medium and also prevents the precipitation of macroscopic crystals. To do this, acids with hydrophilic counterions will preferably be used, for instance hydrochloric acid, acetic acid and
35 citric acid.

These catanionic membranes are stable in physiological media and make it possible especially to retain by adsorption and/or encapsulation and to control the slow diffusion of pharmaceutical or

cosmetic active molecules, or alternatively of cells such as bacteria, for example. They are capable in particular of withstanding high osmotic compressions and even of withstanding the conversion of water into
5 ice, while at the same time remaining in the form of clearly individualized aggregates with rigid and locally flat walls.

A subject of the present invention is also a process for preparing a catanionic membrane in
10 accordance with the invention and as described above, said process being characterized in that it comprises the following steps:

1) a first step of formation of unilamellar vesicles by mixing, in an aqueous solvent of low
15 conductivity:

- a) a cationic surfactant (CS) with OH^- counterions in a molar amount Q_{CS} and
- b) one or more anionic surfactants (AS) in a molar amount Q_{AS} strictly greater than
20 Q_{CS} , and corresponding to equations (1) to (3) below:

$$Q_{\text{AS}} = Q_{\text{AS1}} + Q_{\text{AS2}} \quad (1)$$

$$Q_{\text{AS1}} = Q_{\text{CS}} \quad (2) \text{ and}$$

$$Q_{\text{AS2}} < 2(Q_{\text{CS}}) \quad (3)$$

25 in which:

- Q_{AS1} is the molar amount of an anionic surfactant with H^+ counterions
- Q_{AS2} is the molar amount of an anionic surfactant with H^+ counterions or with
30 metal counterions, said surfactants having a carbon-based chain identical to that of the CS or of the AS with H^+ counterions used in an amount Q_{AS1} ,

said mixture of cationic surfactant and of
35 anionic surfactant being prepared at a temperature above the melting point of the chains of said surfactants;

2) a second step of obtaining flat aggregates formed from only one interdigitated or non-inter-

digitated crystalline molecular bilayer, by cooling the mixture obtained in the first step to a temperature below the melting point of the chains of the surfactants present in the mixture;

5 3) a third step of stabilizing the crystalline molecular bilayers obtained above in the second step, by adding at least one neutral and hydrophobic polymer or a polymer of weakly negative overall electrical charge dissolved in an aqueous
10 solvent of low conductivity, said step being performed at a temperature below the melting point of the chains of the surfactants present in the mixture.

During the first step, the dissolution of the surfactants in the aqueous solvent is preferably
15 performed slowly and may especially take place over variable times ranging from one hour to one week, with minimum mechanical stirring, without heating.

According to a first particular embodiment of this process, and when the anionic surfactants used
20 during the first step consist only of AS with H^+ counterions, this is then referred to as a true catanionic mixture thus comprising only OH^- counterions (provided by the CSs) and H^+ counterions. In this case, the mixture of the ASs and the CSs may optionally be
25 prepared beforehand in powder form before dissolving in the solvent.

According to a second embodiment of this process, and when the excess AS consists of ASs with metal counterions, then the first step of the process
30 in accordance with the invention comprises:

- a first substep in which the CS with OH^- counterions is first mixed with the AS with H^+ counterions in an amount Q_{AS1} equal to Q_{CS} , and then

- a second substep in which the molar amount
35 Q_{AS2} of AS with metal counterions is then added.

In this case, the nature of the metal counterions may be chosen as a function of the property that it is desired to give to the cationic membrane in accordance with the present invention. By way of

example, mention may be made especially of sodium counterions with anticorrosive properties, and also chromium, vanadium and nickel ions, which have electro-chemical corrosion-inhibiting properties on iron alloys.

Once the surfactant solutions are optically homogeneous, the heating step makes it possible to completely melt the chains of the surfactants associated with the pairs of ions formed during the slow dissolution. During this step, the compounds intended to form the bilayers in accordance with the invention are dispersed in the form of vesicles with high electrostatic repulsion. Each vesicle constitutes a microreactor formed from a fluid bilayer (liquid chains), which, after the cooling step, will be converted into a rigid bilayer (gelled chains).

The aqueous solvents used during this process preferably have a conductivity of less than or equal to about 1 MOhm. They are preferably chosen from water and glycerol, and mixtures thereof.

During the first step, the total concentration of surfactants (AS + CS) in the solution is preferably between 0.01% and 3% by weight relative to the total weight of said solution.

Also during the first step, the temperature to which the mixture is heated obviously depends on the nature of the cationic and anionic surfactants used. However, in general, this temperature is generally greater than 30°C and less than 80°C and even more preferentially between 30°C and 70°C. This temperature may be adjusted for each surfactant solution, to 5°C above the temperature of the mixture of surfactants under consideration, which may be detected by an endothermic peak by differential scanning calorimetry (DSC).

This temperature is even more preferentially between 55°C and 70°C. Thus, during the second step, the mixture is cooled to a temperature preferably less than 30°C and even more preferentially to a temperature

of between 20°C and 25°C.

According to one particular embodiment of the process in accordance with the invention, at least one active substance that is absorbed onto the surface of
5 the catanionic membranes and/or encapsulated inside the vesicles (molecular boxes) may also be added to the mixture during the first step. Among these active substances, mention may be made especially of pharmaceutical active principles, active substances for
10 cosmetic purposes and in particular volatile odoriferous molecules, cells such as whole bacteria, and DNA or RNA fragments.

At this moment, a person skilled in the art will preferably take care to select active substances
15 whose electrical charge will be sufficiently low, so as to avoid any destabilization of the catanionic membranes in accordance with the invention.

According to one particular embodiment of the process in accordance with the invention, it is also
20 possible, during the second step, to adjust the pH of the mixture to between 2 and 6 as indicated previously.

According to one advantageous embodiment of the invention and when the catanionic membranes are in the form of hollow polyhedra (molecular boxes), then
25 the process generally includes an additional step of removing the active substances that have not been encapsulated in the polyhedra or that have been adsorbed onto their surface. This removal step may be performed by rinsing, especially using an aqueous
30 solvent identical to the solvent used for the preparation of the membranes, by dialysis or alternatively by filtration.

According to the invention, the volume fraction of polymer added to the mixture during the
35 third step is preferably between one and two times the total mass of the cationic and anionic surfactants in order to have a steric or electrostatic protective layer allowing the addition of salt without destroying the faceted polyhedra of icosahedron type, or the

precipitation of catanionic species forming a three-dimensional crystal.

Finally, a subject of the invention is also the catanionic membranes in accordance with the invention and as described above, for use as medicament for the vectorization of active species or for the retention by adsorption and slow diffusion of volatile molecules.

According to one advantageous embodiment of the invention, the catanionic membrane is in the form of a faceted hollow polyhedron and is used for:

- encapsulating medicaments, for the purpose of vectorizing them,
- encapsulating whole bacteria or DNA or RNA fragments, so as to make them inaccessible to the immune system,
- retaining reagents for chemical reactions taking place inside the polyhedra,
- performing precipitation or crystallization reactions inside the polyhedra, by slow diffusion of reagents inside across the pores of the polyhedra,
- as cosmetic ingredient for the manufacture of creams, obtained by flocculation in the form of bunches of polyhedra, and allowing the efficient diffusion of active molecules after adsorption of the polyhedron onto surfaces of opposite surface electrical potential, for instance the skin.

Besides the preceding arrangements, the invention also comprises other arrangements that will emerge from the description that follows, which refers to an example of preparation of crystalline hollow polyhedra based on cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions and an example of preparation of crystalline hollow polyhedra based on cetyltrimethylammonium hydroxide and myristic acid with H^+ and Li^+ counterions, and also to the attached figures 1 to 4, in which:

- figure 1 is a cryofracture microscopy photograph of crystalline hollow polyhedra based on

cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions before stabilization with a neutral polymer (PEG 20 000) and at a total concentration of surfactants in water of 1% by weight and with a mole
5 fraction AS/(AS + CS) equal to 0.56;

- figure 2 is a cryofracture microscopy photograph of crystalline hollow polyhedra based on cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions after stabilization with a neutral
10 polyethylene glycol polymer (PEG 20 000) and at a total concentration of surfactants in water of 1% by weight and with a mole fraction AS/(AS + CS) equal to 0.56;

- the attached figure 3 shows the X-ray diffraction spectra at 25°C (crystalline structure) and
15 at 65°C (noncrystalline structure) of catanionic mixtures of CTAOH and of myristic acid with H^+ counterions, on which the diffusion intensity (in cm^{-1}) is expressed as a function of the wave vector Q (in \AA^{-1});

- figure 4 is a cryofracture microscopy
20 photograph of crystalline hollow polyhedra based on cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions in equimolar amounts, and of an excess of myristic acid with Li^+ counterions.

It should be clearly understood, however,
25 that these examples are given solely as illustrations of the subject of the invention, of which they do not in any way constitute a limitation.

EXAMPLE 1: Preparation of crystalline hollow polyhedra
30 **based on cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions**

This example illustrates one of the two variants of the process for preparing the bilayers in accordance with the invention, namely that in which the
35 CS with OH^- counterions and the AS with H^+ counterions in excess are mixed together directly to obtain a true catanionic mixture, without passing via a prior substep of equimolar premixing of AS and CS.

To do this, 0.023 g of cetyltrimethylammonium

hydroxide (CTAOH) and 0.022 g of myristic acid with H^+ counterions are mixed together, in the form of freeze-dried powders. A mixture whose mole fraction of myristic acid with H^+ counterions/(myristic acid with H^+ counterions + CTAOH) is equal to 0.56 is thus obtained. 4.46 g of Millipore water ($< 1 \text{ MOhm/cm}$) is then added to the powder mixture thus obtained to form a solution with a total weight concentration of surfactants equal to 1% and a mole fraction AS/(AS + CS) equal to 0.56. The compounds are left to dissolve at room temperature with gentle stirring for one week. After the surfactants have fully dissolved (disappearance of the solid grains of myristic acid), the solution is heated to a uniform temperature and slightly higher than 65°C , which corresponds to the melting point of the chains of the surfactants, for one minute. The solution is then allowed to cool to room temperature.

Separately, a solution containing 1.5% by weight of polyethylene glycol sold under the name PEG 20 000 (neutral polymer) is prepared. The surfactant solution is mixed at room temperature, on a volume-for-volume basis, with the polymer solution with gentle stirring. The final dispersion obtained may be concentrated by simple filtration on account of the conjunction of the crystallization of the chains, the size of the objects, the low viscosity of the solution and the low osmotic pressure ($< 1000 \text{ Pa}$) of the final dispersion obtained.

The solution obtained before concentration is slightly scattering and bluish and contains a dispersion of hollow micron-sized faceted objects that may be detected by their characteristic scattering decreasing with the square of the scattering angle at small angles (light or neutrons), associated with the presence of a fine Bragg peak ($< 0.002 \text{ nm}$) located between 0.150 nm^{-1} and 0.156 nm^{-1} by large-angle X-ray scattering.

Moreover, the stability properties of the membrane according to the invention in the presence of

saline solutions up to isotonic osmolarity were demonstrated by mixing with a concentrated sodium chloride solution until a final saline concentration of 0.15 M was obtained.

5 The icosahedric boxes thus obtained were observed directly by cryofracture microscopy. The images obtained are given in the attached figures 1 and 2, taken before and after stabilization with the neutral polymer. The crystalline nature of the faces
10 was moreover confirmed by X-ray scattering on samples of identical chemical nature but more concentrated (band at $q = 1.52 \pm 0.01 \text{ \AA}^{-1}$ as is seen in the attached figure 3 that shows the scattering intensity (in cm^{-1}) as a function of the wave vector Q (in \AA^{-1}) at a
15 temperature of 65°C (noncrystalline structure) and at a temperature of 25°C (crystalline structure).

EXAMPLE 2: Preparation of crystalline hollow polyhedra based on cetyltrimethylammonium hydroxide and myristic acid with H^+ counterions and an excess of myristic acid with Li^+ counterions
20

This example illustrates the second variant of the process in accordance with the invention, i.e. the preparation of hollow polyhedra in which the excess
25 of anionic surfactant is composed of an anionic surfactant with lithium counterions, which imposes the implementation of the first step of the process in two substeps as explicitly described previously.

Just as in example 1, 0.0230 g of freeze-dried CTAOH and 0.0174 g of myristic acid with H^+
30 counterions are first mixed together and an amount of Millipore water ($> 1 \text{ MOhm/cm}$) sufficient to form a solution containing 1% by weight of total surfactants is then added. In this solution, the molar amount Q_c of CTAOH is identical to the molar amount Q_{A1} of myristic
35 acid with H^+ counterions. The surfactant mixture is then allowed to dissolve by slow stirring at room temperature until a solution that no longer contains any heterogeneities visible to the naked eye and

corresponding to myristic acid crystals is obtained. The solution is then heated to a temperature above 50°C for 1 minute; the solution should be transparent and should not contain any aggregates. It is allowed to
5 cool to room temperature. 0.0051 g of powdered lithium myristate is then added to this solution. A solution of catanionic surfactants is then obtained, in which the mole fraction of myristate ($H^+ + Li^+$)/(myristate ($H^+ + Li^+$) + CTAOH) is equal to 0.56. The mixture is stirred
10 at room temperature for one week until the lithium myristate has fully dissolved. The catanionic solution is then heated to a temperature above 65°C for 1 minute, and the solution is then allowed to cool to room temperature. The objects obtained by this method
15 have the same structure as those of example 1 above and are shown in the attached figure 4 obtained by cryo-fracture microscopy.

Separately, a solution containing 1.5% by weight of neutral polymer: polyethylene glycol (PEG
20 20 000) is prepared. The catanionic solution is mixed at room temperature, volume-for-volume, with the neutral polymer solution with gentle stirring. As in example 1 above, the final dispersion obtained may be concentrated by simple filtration.

25 The structural signature (not shown) is similar to that of example 1, both by cryofracture and by X-ray.